



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

FOREIGN AND INSULAR STATISTICAL REPORTS OF COUNTRIES AND CITIES—UNTABULATED.

AUSTRALIA—*New South Wales*—*New Castle*.—Month of October, 1905. Census population, 68,017. Total number of deaths 40, including diphtheria 1, and 5 from tuberculosis.

BRITISH GUIANA—*Demerara*—*Georgetown*.—Month of October, 1905. Estimated population, 36,567. Total number of deaths 236, including 24 from tuberculosis.

CANADA—*British Columbia*—*Victoria*.—Month of October, 1905. Estimated population, 21,000. Total number of deaths 25, including enteric fever 2, and 1 from tuberculosis.

CUBA—*Habana*.—Month of October, 1905. Estimated population, 275,000. Total number of deaths, 513, including diphtheria 1, enteric fever 3, measles 2, whooping cough, 1, yellow fever 2, and 113 from tuberculosis.

FRANCE—*St. Etienne*.—Two weeks ended November 15, 1905. Estimated population, 146,671. Total number of deaths 121, including diphtheria 1, enteric fever 2, whooping cough 2, and 10 from tuberculosis.

GERMANY—*Weimar*.—Month of October, 1905. Estimated population, 30,829. Total number of deaths 45, including diphtheria 1, enteric fever 2, and 4 from tuberculosis.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 76 great towns in England and Wales during the week ended November 18, 1905, correspond to an annual rate of 15.5 per 1,000 of the aggregate population, which is estimated at 15,609,377.

Bradford.—Two weeks ended November 18, 1905. Estimated population, 286,799. Total number of deaths, 169, including diphtheria 4, scarlet fever 3, and 12 from tuberculosis.

London.—One thousand five hundred and fifty-two deaths were registered during the week, including measles 54, scarlet fever 8, diphtheria 19, whooping cough 20, enteric fever 8, and 13 from diarrhea. The deaths from all causes correspond to an annual rate of 17.3 per 1,000. In Greater London 2,050 deaths were registered. In the "outer ring" the deaths included 3 from diphtheria, 9 from measles, 5 from whooping cough, and 4 from diarrhea.

Ireland.—The average annual death rate represented by the deaths registered during the week ended November 18, 1905, in the 21 principal town districts of Ireland was 19.8 per 1,000 of the population, which is estimated at 1,093,959. The lowest rate was recorded in Waterford, viz, 3.9, and the highest in Clonmel, viz, 41 per 1,000. In Dublin and suburbs 158 deaths were registered, including diphtheria 1, scarlet fever 1, whooping cough 1, and 38 from tuberculosis.

Scotland.—The deaths registered in 8 principal towns during the week ended November 18, 1905, correspond to an annual rate of 17.5

per 1,000 of the population, which is estimated at 1,749,917. The lowest rate of mortality was recorded in Paisley and Leith, viz, 12.8, and the highest in Greenock, viz, 22.3 per 1,000. The aggregate number of deaths registered from all causes was 588, including measles 34, diphtheria 4, enteric fever 2, scarlet fever 3, and 2 from whooping cough.

JAMAICA—*Kingston*.—Month of October, 1905. Estimated population, 52,475. Total number of deaths, 114, including 11 from phthisis pulmonalis.

JAPAN—*Formosa*.—Month of October, 1905. Estimated population, 2,922,585. Total number of deaths not reported. One death from diphtheria, 2 deaths from enteric fever, and 1 death from plague reported.

NAGASAKI.—Ten days ended October 31, 1905. Estimated population, 159,041. Total number of deaths not reported. No deaths from contagious diseases reported.

Ten days ended November 10, 1905. Estimated population, 159,041. Total number of deaths not reported. One death from enteric fever reported.

MALTA.—Two weeks ended November 11, 1905. Estimated population, 202,134. Total number of deaths, 206, including diphtheria 2, and 1 from enteric fever.

SPAIN—*Barcelona*.—Ten days ended November 20, 1905. Estimated population, 600,000. Total number of deaths, 409, including diphtheria 5, enteric fever 19, measles 3, scarlet fever 1, whooping cough 1, smallpox 6, and 30 from tuberculosis.

CADIZ.—Month of October, 1905. Estimated population, 69,382. Total number of deaths 122, including enteric fever 1, measles 1, and 29 from tuberculosis.

TURKEY—*Constantinople*.—Two weeks ended November 19, 1905. Estimated population, 800,000. Total number of deaths 424, including diphtheria 1, enteric fever 16, measles 2, scarlet fever 2, and 7 from smallpox.

WEST INDIES—*Curaçao*.—Two weeks ended November 24, 1905. Estimated population, 31,600. Total number of deaths 19. No contagious diseases.

ST. THOMAS.—Two weeks ended November 24, 1905. Estimated population, 12,019. Total number of deaths 10, including 2 from tuberculosis.